

15
16. (New) Detection method, without labelling, of a molecular recognition reaction according to Claim 15, which is a hybridization reaction of nucleic acids consisting of the following steps:

- fixation of the first nucleic acid molecule on a solid support,
- contacting of the first nucleic acid molecule fixed on the solid support with a solution to be tested suspected of containing the second nucleic acid molecule, this latter being capable of being hybridized with said first molecule, the contacting being carried out under conditions favourable for said hybridization,

- washing of the solid support to isolate a detection sample formed from said first molecule fixed on the support and possibly said second molecule hybridized on said first molecule, and

- measuring the absorption of the sample by a photothermal method.

16
17. (New) Method according to Claim 14 in which the photothermal method is a thermal lens method.

17
18. (New) Method according to Claim 15 in which the photothermal method is a thermal lens method.

18
19. (New) Method according to Claim 15 in which the photothermal method is a method of photothermal deflection in which the sample is illuminated by a pump beam and the absorption of the pump beam by a sample is detected by the refraction or the reflection of a probe beam.

19
20. (New) Method according to Claim 15 in which the photothermal method is a method of photothermal deflection in which the sample is illuminated by a pump beam and the absorption of the pump beam by a sample is detected by the refraction or the reflection of a probe beam.

20

21. (New) Method according to Claim ¹⁸~~19~~ in which the probe and pump beams cross each other.

21

22. (New) Method according to Claim ¹⁹~~20~~ in which the probe and pump beams cross each other.

22

23. (New) Method according to Claim ¹⁸~~19~~ in which the probe and pump beams are in a transverse configuration or in an approximately collinear configuration.

23

24. (New) Method according to Claim ¹⁹~~20~~ in which the probe and pump beams are in a transverse configuration or in an approximately collinear configuration.

24

25. (New) Method according to Claim ¹⁸~~19~~ in which the pump beam is chosen from a pulsed laser, a continuous intensity modulated laser or polychromatic light.

25

26. (New) Method according to Claim ¹⁸~~19~~ in which the refraction or the reflection of the probe beam is detected by means of multielement photodiode or by means of a simple photodiode receiving only part of the probe beam.

26

27. (New) Method according to Claim ²⁴~~25~~ in which the refraction or the reflection of the probe beam is detected by means of multielement photodiode or by means of a simple photodiode receiving only part of the probe beam.

27

28. (New) Method according to Claim ¹⁸~~19~~ in which the pump beam is a beam from a laser chosen from a continuous argon laser at 275 nm, a quadrupled YAG laser with a wavelength of 266 nm or a polychromatic light.

28

29. (New) Method according to Claim ¹⁸~~19~~ in which the probe beam has a wavelength that is not absorbed by the substrate nor the present molecules.

29

30. (New) Method according to Claim ¹⁶~~17~~ in which an incident beam is used, said beam being a beam from a laser chosen from a continuous argon laser at 275 nm, a quadrupled YAG laser with a wavelength of 266 nm or polychromatic light.

30

31. (New) Method according to Claim ¹⁵~~16~~ comprising in addition a step for comparing the measurement of absorption of the sample with that of a control sample.

31

32. (New) Use of a method according to Claim ¹⁴~~15~~ for a test, a diagnosis or a detection of hybridization of nucleic acids.

32

33. (New) Use of a method according to Claim 16 for a test, a diagnosis or a detection of hybridization of nucleic acids.

33

34. (New) Method according to Claim ¹⁹~~20~~ in which the pump beam is chosen from a pulsed laser, a continuous intensity modulated laser or polychromatic light.

34

35. (New) Method according to Claim ¹⁹~~20~~ in which the refraction or the reflection of the probe beam is detected by means of multielement photodiode or by means of a simple photodiode receiving only part of the probe beam.

35

36. (New) Method according to Claim ³³~~34~~ in which the pump beam is chosen from a pulsed laser, a continuous intensity modulated laser or polychromatic light.

36

37. (New) Method according to Claim ¹⁹~~20~~ in which the pump beam is a beam from a laser chosen from a continuous argon laser at 275 nm, a quadrupled YAG laser with a wavelength of 266 nm or a polychromatic light.

37

38. (New) Method according to Claim ¹⁹~~20~~ in which the probe beam has a wavelength that is not absorbed by the substrate nor the present molecules.

38

39. (New) Use of a method according to Claim ²⁴~~25~~ for a test, a diagnosis or a detection of hybridization of nucleic acids.